

A new approach to hyperbolic inverse problems.

G.Eskin, Department of Mathematics, UCLA,
Los Angeles, CA 90095-1555, USA. E-mail: eskin@math.ucla.edu

February 1, 2008

Abstract

We present a modification of the BC-method in the inverse hyperbolic problems. The main novelty is the study of the restrictions of the solutions to the characteristic surfaces instead of the fixed time hyperplanes. The main result is that the time-dependent Dirichlet-to-Neumann operator prescribed on a part of the boundary uniquely determines the coefficients of the self-adjoint hyperbolic operator up to a diffeomorphism and a gauge transformation. In this paper we prove the crucial local step. The global step of the proof will be presented in the forthcoming paper.

1 Introduction.

Consider a hyperbolic equation of the form

$$Lu \stackrel{\text{def}}{=} \frac{\partial^2 u}{\partial t^2} + \sum_{j,k=1}^n \frac{1}{\sqrt{g(x)}} \left(-i \frac{\partial}{\partial x_j} + A_j(x) \right) \sqrt{g(x)} g^{jk}(x) \left(-i \frac{\partial}{\partial x_k} + A_k(x) \right) u + V(x)u = 0 \quad (1.1)$$

in $\Omega \times (0, T_0)$, where Ω is a smooth bounded domain in $\mathbf{R}^n$, all coefficients in (1.1) are $C^\infty(\overline{\Omega})$ functions, $\|g^{jk}(x)\|^{-1}$ is the metric tensor in $\overline{\Omega}$, $g(x) = \det \|g^{jk}\|^{-1}$. We assume that

$$(1.2) \quad u(x, 0) = u_t(x, 0) = 0 \quad \text{in } \Omega, \quad u|_{\partial\Omega \times (0, T_0)} = f(x, t).$$

We shall study the inverse boundary problem with boundary data given on the part of the boundary. Let Γ_0 be an open subset of $\partial\Omega$ (in particular, $\Gamma_0 = \partial\Omega$) and let the D-to-N (Dirichlet-to-Neumann) operator

$$(1.3) \quad \Lambda f = \sum_{j,k=1}^n g^{jk}(x) \left(\frac{\partial u}{\partial x_j} + iA_j(x)u \right) \nu_k \left(\sum_{p,r=1}^n g^{pr}(x) \nu_p \nu_r \right)^{-\frac{1}{2}} \Big|_{\Gamma_0 \times (0, T_0)}$$

be given for all f with the supports in $\Gamma_0 \times (0, T_0]$. Here $\nu = (\nu_1, \dots, \nu_n)$ is the unit exterior normal to $\partial\Omega$ with respect to the Euclidian metric. If $F(x) = 0$ is the equation of $\partial\Omega$ in some neighborhood then (1.3) has the following form in this neighborhood:

$$(1.4) \quad \Lambda f = \sum_{j,k=1}^n g^{jk}(x) \left(\frac{\partial u}{\partial x_j} + iA_j(x)u \right) F_{x_j}(x) \left(\sum_{p,r=1}^n g^{pr}(x) F_{x_p} F_{x_r} \right)^{-\frac{1}{2}} \Big|_{F(x)=0, 0 < t < T_0}.$$

Let

$$(1.5) \quad y = y(x)$$

be a C^∞ diffeomorphism of $\overline{\Omega}$ onto $\overline{\Omega_0}$ such that $\Gamma_0 \subset \partial\Omega_0$, the Jacobian $\det \frac{\mathcal{D}y}{\mathcal{D}x} \neq 0$ in $\overline{\Omega}$ and

$$(1.6) \quad y = x \quad \text{on} \quad \Gamma_0.$$

If we change variables $y = y(x)$ in (1.1) we get an equation in Ω_0 of the same form as (1.1):

$$(1.7) \quad \frac{\partial^2 v}{\partial t^2} + \sum_{j,k=1}^n \frac{1}{\sqrt{g_0(y)}} \left(-i \frac{\partial}{\partial y_j} + A_j^{(0)}(y) \right) \sqrt{g_0(y)} g_0^{jk}(y) \left(-i \frac{\partial}{\partial y_k} + A_k^{(0)}(y) \right) v(y, t) + V^{(0)}(y) v(y, t) = 0,$$

where $v(y(x), t) = u(x, t)$,

$$(1.8) \quad \|g_0^{jk}(y(x))\| = \left(\frac{\mathcal{D}y}{\mathcal{D}x} \right) \|g^{jk}(x)\| \left(\frac{\mathcal{D}y}{\mathcal{D}x} \right)^T,$$

$$(1.9) \quad g_0(y) = \det \|g_0^{jk}(y)\|^{-1}, \quad V^{(0)}(y(x)) = V(x)$$

and

$$(1.10) \quad y^*(A^{(0)}) = A,$$

where

$$(1.11) \quad A = \sum_{j=1}^n A_j(x) dx_j, \quad A^{(0)} = \sum_{j=1}^n A_j^{(0)}(y) dy_j.$$

The D-to-N operator $\Lambda^{(0)}$ corresponding to (1.7) has the form
(1.12)

$$\Lambda^{(0)} f = \sum_{j,k=1}^n g_0^{jk}(y) \left(\frac{\partial v}{\partial y_j} + i A_j^{(0)}(y) v \right) \nu_k \left(\sum_{p,r=1}^n g_0^{pr}(y) \nu_p \nu_r \right)^{-\frac{1}{2}} \Big|_{\Gamma_0 \times (0, T_0)},$$

where $v(y, t) \Big|_{\Gamma_0 \times (0, T_0)} = f(y, t)$ since (1.5) is the identity on Γ_0 .

It follows from (1.4) and (1.6) that

$$(1.13) \quad \Lambda^{(0)} f = \Lambda f, \quad \forall f, \quad \text{supp } f \subset \Gamma_0 \times (0, T_0].$$

Denote by $G_0(\overline{\Omega})$ the group of $C^\infty(\overline{\Omega})$ functions such that $c(x) \neq 0$ on $\overline{\Omega}$ and $c(x) = 1$ on Γ_0 . We say that $A(x) = (A_1(x), \dots, A_n(x))$ and $A^{(1)} = (A_1^{(1)}(x), \dots, A_n^{(1)}(x))$ are gauge equivalent in Ω if there exists $c(x) \in G_0(\overline{\Omega})$ such that

$$A_j^{(1)}(x) = A_j(x) - i c^{-1}(x) \frac{\partial c}{\partial x_j}, \quad 1 \leq j \leq n.$$

If $u^{(1)}(x, t) = c^{-1}(x) u(x, t)$ where $u(x, t)$ is the solution of (1.1) then $L^{(1)} u^{(1)} = 0$ where $L^{(1)}$ is the same as L with $A(x)$ replaced by $A^{(1)}(x)$. We shall write for the brevity that $c \circ L^{(1)} = L$.

Denote

$$T_* = \sup_{x \in \overline{\Omega}} d(x, \Gamma_0),$$

where $d(x, \Gamma_0)$ is the distance in $\overline{\Omega}$ with respect to metric tensor $\|g^{jk}\|^{-1}$ from $x \in \Omega$ to Γ_0 .

Theorem 1.1. *Suppose $T_0 > 2T_*$ and L and L_0 are operators of the form (1.1) and (1.7) in Ω and Ω_0 respectively, where $A_j(x), V(x)$ and $A_j^{(0)}, V^{(0)}$ are real valued, $1 \leq j \leq n$. Let Λ be the D-to-N operator corresponding to (1.1). If $\Lambda^{(0)}$ is the D-to-N corresponding to (1.7) and $\Lambda = \Lambda^{(0)}$ on $\Gamma_0 \times (0, T_0)$ then*

there exists a diffeomorphism (1.5) and a gauge transformation $c(x) \in G_0(\overline{\Omega})$ such that

$$c \circ y^{-1} \circ L^{(0)} = L.$$

The first result in this direction was obtained in [I]. The most general results were obtained by the BC-method (see [B1], [B2], [K], [KK] and [KKL]).

An important class of inverse problems with the data given on a part of the boundary is the inverse problems in domains with obstacles. In this case Ω is a ball B_R with removed subdomains $\Omega_1, \dots, \Omega_m$, called obstacles. The D-to-N operator Λ is given on $\partial B_R \times (0, T_0)$ and the zero Dirichlet boundary conditions are satisfied on $\partial \Omega_j \times (0, T_0)$, $j = 1, \dots, m$. In [E1] such inverse problems were considered in the connection with the Aharonov-Bohm effect assuming that B_R contains one or several convex obstacles.

In the present paper we developed a modification of the BC-method. The main novelty is the study of the restrictions of the solutions to the characteristic surface instead of the restrictions to the hyperplane $t = \text{const}$ as in BC-method. The proof of Theorem 1.1 consists of two steps. In the first step we prove that knowing Λ in the neighborhood of $\Gamma \times (0, T)$, where $\Gamma \subset \Gamma_0$, $0 < T < T_0$, one can recover the coefficients of the equation (1.1) in some neighborhood of Γ up to a diffeomorphism and a gauge transformation. This will be done in §2. In §3 we prove some lemmas used in §2. The global step of the proof of Theorem 1.1 will be given in the forthcoming paper. A generalization to the case of Yang-Mills potentials is considered in [E4].

2 The local step.

Let Γ be an open subset of Γ_0 and let U_0 be a neighborhood of Γ . Let (x', x_n) be a system of coordinates in U_0 such that the equation of Γ_0 is $x_n = 0$ and $x' = (x_1, \dots, x_{n-1})$ are coordinates on $\Gamma_0 \cap U_0$.

We introduce semi-geodesic coordinates for L . Denote by $\varphi_n(x)$ the solution of the equation (c.f.[E3])

$$(2.1) \quad \sum_{j,k=1}^n g^{jk}(x) \frac{\partial \varphi_n}{\partial x_j} \frac{\partial \varphi_n}{\partial x_k} = 1, \quad 0 \leq x_n < \delta,$$

$$\varphi_n(x', 0) = 0.$$

Also denote by $\varphi_p(x)$, $1 \leq p \leq n-1$, the solutions of the equations

$$(2.2) \quad \sum_{j,k=1}^n g^{jk}(x) \frac{\partial \varphi_n}{\partial x_j} \frac{\partial \varphi_p}{\partial x_k} = 0,$$

$$\varphi_p(x', 0) = x_p, \quad 1 \leq p \leq n-1.$$

We have that $y = \varphi(x) = (\varphi_1(x), \dots, \varphi_n(x))$ exists for $0 \leq x_n \leq \delta$ when δ is small and the Jacobian

$$J(x) = \left| \frac{\mathcal{D}\varphi}{\mathcal{D}x} \right| \neq 0 \quad \text{in } \bar{\Gamma} \times [0, \delta].$$

Note that $\varphi(x', 0) = x'$. Changing variables $y = \varphi(x)$ in (1.1) we get

$$\hat{L}\hat{u} = 0,$$

where $\hat{u}(y, t) = u(x, t)$ and

$$(2.3) \quad \hat{L} = \frac{\partial^2}{\partial t^2} + \sum_{j,k=1}^n \frac{1}{\sqrt{\hat{g}(y)}} \left(-i \frac{\partial}{\partial y_j} + \hat{A}_j(y) \right) \sqrt{\hat{g}(y)} \hat{g}^{jk}(y) \left(-i \frac{\partial}{\partial y_k} + \hat{A}_k(y) \right) + \hat{V}(y),$$

$$(2.4) \quad \hat{g}^{jk}(\varphi(x)) = \sum_{p,r=1}^n g^{pr}(x) \frac{\partial \varphi_j}{\partial x^p} \frac{\partial \varphi_k}{\partial x^r}, \quad \hat{g}(y) = \det \|\hat{g}^{jk}(y)\|^{-1}, \quad \hat{V}(\varphi(x)) = V(x)$$

and

$$A_k(x) = \sum_{j=1}^n \hat{A}_j(\varphi(x)) \frac{\partial \varphi_j(x)}{\partial x_k}.$$

It follows from (2.1), (2.2) that

$$(2.5) \quad \hat{g}^{nn}(y) = 1, \quad \hat{g}^{nj} = \hat{g}^{jn} = 0, \quad 1 \leq j \leq n-1.$$

Let

$$(2.6) \quad A'_j(y) = -\frac{i}{2}(\sqrt{\hat{g}})^{-1} \frac{\partial \sqrt{\hat{g}}}{\partial y_j} = -\frac{i \hat{g}_{y_j}}{4 \hat{g}}, \quad 1 \leq j \leq n.$$

Then we can rewrite (2.3) in the form

$$(2.7) \quad \hat{L}\hat{u} = \frac{\partial^2 \hat{u}}{\partial t^2} + \left(-i \frac{\partial}{\partial y_n} + \hat{A}_n(y) + A'_n(y) \right)^2 \hat{u} \\ + \sum_{j,k=1}^{n-1} \left(-i \frac{\partial}{\partial y_j} + \hat{A}_j + A'_j(y) \right) \hat{g}^{jk} \left(-i \frac{\partial}{\partial y_k} + \hat{A}_k + A'_k \right) \hat{u} + V_1(y) \hat{u} = 0,$$

where

$$(2.8) \quad V_1 = (A'_n)^2 + i \frac{\partial A'_n}{\partial y_n} + \sum_{j,k=1}^{n-1} \left(\hat{g}^{jk} A'_j A'_k + i \frac{\partial}{\partial y_j} (\hat{g}^{jk} A'_k) \right) + \hat{V}(y).$$

Note that $\hat{u}(y, 0) = \hat{u}_t(y, 0) = 0$, $\hat{u}|_{\Gamma \times (0, T)} = f(y', t)$ since $\varphi(x', 0) = x'$.

Denote by $\hat{\Lambda}$ the D-to-N operator corresponding to $\hat{L}$. We have

$$(2.9) \quad \hat{\Lambda}f = \left(\frac{\partial \hat{u}(y', y_n, t)}{\partial y_n} + i \hat{A}_n(y', y_n) \hat{u}(y', y_n, t) \right) \Big|_{\Gamma \times (0, T_0)}.$$

Note that

$$(2.10) \quad \Lambda f = \hat{\Lambda}f \quad \text{on } \Gamma \times (0, T_0),$$

where $\text{supp } f \subset \Gamma \times (0, T_0)$. It will be shown in Remark 2.2 that the D-to-N operator $\hat{\Lambda}$ on $\Gamma \times (0, T_0)$ determines the restriction of the metric tensor $\|\hat{g}^{jk}\|^{-1}$ to Γ and, in particular,

$$(2.11) \quad \hat{g}(y', 0) \quad \text{and} \quad \frac{\partial \hat{g}}{\partial y_n}(y', 0).$$

Make the transformation $\hat{u} = (\hat{g}(y', y_n))^{-\frac{1}{4}} u'$. Then $L'u' = 0$ in $U_0 \times (0, T_0)$ where L' is similar to (2.7) with $\hat{A}_j + A'_j$ replaced by $\hat{A}_j$, $1 \leq j \leq n$. Denote by Λ' the D-to-N operator corresponding to L' , i.e.

$$(2.12) \quad \Lambda' f' = \left(\frac{\partial u'}{\partial y_n} + i \hat{A}_n u' \right) \Big|_{\Gamma \times (0, T)} \quad \text{where} \quad f' = u' \Big|_{\Gamma \times (0, T)}$$

We have

$$(2.13) \quad \Lambda' f' = \left(\frac{\partial}{\partial y_n} + i \hat{A}_n \right) ((\hat{g})^{\frac{1}{4}} \hat{u}) \Big|_{\Gamma \times (0, T)} = \hat{g}^{\frac{1}{4}}(y', 0) \hat{\Lambda}f + \frac{\hat{g}_{y_n}(y', 0)}{4\hat{g}(y', 0)} f'(y', 0, t).$$

It follows from (2.11) that $\hat{\Lambda}$ determines Λ' on $\Gamma \times (0, T_0)$.

Let $\hat{\psi} \in C^\infty(\overline{U_0})$, $\hat{\psi}(y', 0) = 0$. Then $c(y', y_n) = e^{i\hat{\psi}} \in G_0(\overline{U_0})$. We say that $A_j^{(1)}(y)$ and $\hat{A}_j(y)$ are gauge equivalent in $\overline{U_0}$ if

$$(2.14) \quad A_j^{(1)} = \hat{A}_j + \frac{\partial \hat{\psi}}{\partial y_j} \quad \text{in } \overline{U_0}, \quad 1 \leq j \leq n.$$

We shall choose $\hat{\psi}(y)$ such that $A_n^{(1)} = 0$. Denote $u' = e^{i\hat{\psi}} \hat{u}_1$. Then $\hat{u}_1$ satisfies the equation

$$L_1 \hat{u}_1 = 0,$$

where

$$(2.15) \quad L_1 \hat{u}_1 = \frac{\partial^2 \hat{u}_1}{\partial t^2} - \frac{\partial^2 \hat{u}_1}{\partial y_n^2} + \sum_{j,k=1}^{n-1} \left(-i \frac{\partial}{\partial y_j} + A_j^{(1)}(y) \right) \hat{g}^{jk} \left(-i \frac{\partial}{\partial y_k} + A_k^{(1)}(y) \right) \hat{u}_1 + V_1(y) \hat{u}_1 = 0,$$

i.e. L_1 is obtained from L' by replacing $\hat{A}_j$ by $A_j^{(1)}$. Note that the D-to-N operator $\Lambda^{(1)}$ corresponding to L_1 has the form:

$$(2.16) \quad \Lambda^{(1)} f = \frac{\partial \hat{u}_1}{\partial y_n} \Big|_{y_n=0, 0 < t < T_0},$$

since $A_n^{(1)} = 0$. Note that

$$(2.17) \quad \Lambda^{(1)} f = \Lambda' f,$$

where Λ' is the D-to-N corresponding to L' . Denote by L_1^* the formally adjoint operator to L_1 . Then L_1^* has the same form as (2.15) with $A_j^{(1)}, V_1$ replaced by $\overline{A_j^{(1)}}, \overline{V_1}$. Let $\Lambda_*^{(1)}$ be the D-to-N operator corresponding to L_1^* . Note that $\Lambda^{(1)}$ determines $\Lambda_*^{(1)}$. Indeed let $\Lambda^{(1)*}$ be the adjoint operator to $\Lambda^{(1)}$. Then $\Lambda_*^{(1)} f$ is obtained from $\Lambda^{(1)*}$ by changing t to $T - t$ and $f(y', t)$ to $f(y', T - t)$ (c.f. [KL1]). Note that L_1 is self-adjoint if L is self-adjoint. In this case $\Lambda_*^{(1)} = \Lambda^{(1)}$.

Denote $\Delta_{1s_0} = \Gamma \times (s_0, T]$, where $0 \leq s_0 < T$. Let $D(\Delta_{1s_0})$ be the forward domain of influence of Δ_{1s_0} in the half-space $y_n \geq 0$. Denote $\Gamma^{(1)} = \{y' : (y', y_n, t) \in D(\Delta_{1s_0}), y_n = 0, t = T\}$ and $\Delta_{2s_0} = \Gamma^{(1)} \times (s_0, T]$. Let

$D(\Delta_{2s_0})$ be the domain of influence of Δ_{2s_0} in $y_n \geq 0$. Denote by Y_{js_0} the intersection of $D(\Delta_{js_0})$ with the plane $T - t - y_n = 0$ and by X_{js_0} the part of $D(\Delta_{js_0})$ below Y_{js_0} , $j = 1, 2$. Let $Z_{js_0} = \partial X_{js_0} \setminus (Y_{js_0} \cup \{y_n = 0\})$. Also denote $\gamma_{js_0} = \partial Y_{js_0} \cap \{y_n = 0\}$, $1 \leq j \leq 2$. Note that $\gamma_{10} = \Gamma^{(1)}$. Denote by R_{s_0} the following subset of Y_{2s_0} : $\{(s, y') : s_0 \leq s \leq T, y' \in \Gamma^{(1)}\}$, $s = t - y_n$, $T - y_n - t = 0$. We shall choose T such that $D(\Delta_{20})$ does not intersect $\partial\Omega \times [0, T]$ when $y_n > 0$. We also assume that the semigeodesic coordinates are defined in $D(\Delta_{20})$ for $t \leq T$ and that $D(\Delta_{20}) \cap \{y_n = 0\} \subset \Gamma_0 \times [0, T]$ for $t \leq T$.

The main result of this section is the following lemma:

Lemma 2.1. *Let L_1 be self-adjoint. Then the D -to- N operator $\Lambda^{(1)}$ on Δ_{20} determines all coefficients of L_1 (see (2.15)) in $\mathcal{D}_{\frac{T}{2}} = \Gamma \times [0, \frac{T}{2}]$.*

For the most part of the proof of Lemma 2.1 we will not need L_1 to be self-adjoint. We shall use the self-adjointness only in the proof of Lemma 2.4.

Consider the identity

$$(2.18) \quad 0 = (L_1 \hat{u}_1, \frac{\partial \hat{v}_1}{\partial t}) + (\frac{\partial \hat{u}_1}{\partial t}, L_1^* \hat{v}_1),$$

where

$$(2.19) \quad (\hat{u}, \hat{v}) = \int_{X_{20}} \hat{u}(y, t) \overline{\hat{v}(y, t)} dy dt,$$

L_1 is the same as in (2.15), $\hat{u}_1(y', 0, t) = f(y', t)$, $\hat{v}_1(y', 0, t) = g(y', t)$, $\hat{u} = \hat{u}_t = \hat{v} = \hat{v}_t = 0$ for $t = 0, y_n > 0$, $L_1 \hat{u}_1 = 0$, $L_1^* \hat{v}_1 = 0$, $\text{supp } f(y', t) \subset \Delta_{20}$, $\text{supp } g(y', t) \subset \Delta_{20}$. We have:

$$(2.20) \quad \begin{aligned} \int_{X_{20}} (\hat{u}_{1tt} \overline{\hat{v}_{1t}} + \hat{u}_{1t} \overline{\hat{v}_{1tt}}) dy dt &= \int_{X_{20}} \frac{\partial}{\partial t} (\hat{u}_{1t} \overline{\hat{v}_{1t}}) dy dt \\ &= \int_{Y_{20}} \hat{u}_{1t} \overline{\hat{v}_{1t}} dy - \int_{Z_{20}} \hat{u}_{1t} \overline{\hat{v}_{1t}} dy \end{aligned}$$

Since $\hat{u}_1$ and $\hat{v}_1$ have zero Cauchy data for $t = 0$, $y_n > 0$, we get that $\hat{u}_1$, $\hat{v}_1$ are equal to zero on Z_{20} . Integrating by parts in y_n we obtain:

$$(2.21) \quad \begin{aligned} - \int_{X_{20}} (\hat{u}_{1y_n^2} \overline{\hat{v}_{1t}} + \hat{u}_{1t} \overline{\hat{v}_{1y_n^2}}) dy' dy_n dt &= \int_{X_{20}} (\hat{u}_{1y_n} \overline{\hat{v}_{1y_n t}} + \hat{u}_{1y_n t} \overline{\hat{v}_{1y_n}}) dy' dy_n dt \\ &- \int_{Y_{20}} (\hat{u}_{1y_n} \overline{\hat{v}_{1t}} + \hat{u}_{1t} \overline{\hat{v}_{1y_n}}) dy' dt + \int_{\Delta_{20}} (\hat{u}_{1y_n} \overline{\hat{v}_{1t}} + \hat{u}_{1t} \overline{\hat{v}_{1y_n}}) dy' dt. \end{aligned}$$

Note that

$$\hat{u}_{1y_n} \overline{\hat{v}_{1y_n t}} + \hat{u}_{1y_n t} \overline{\hat{v}_{1y_n}} = \frac{\partial}{\partial t} (\hat{u}_{1y_n} \overline{\hat{v}_{1y_n}}).$$

Analogously, integrating by parts in $y' = (y_1, \dots, y_{n-1})$ we get

$$(2.22) \quad \int_{X_{20}} \left[\sum_{j,k=1}^{n-1} \left(-i \frac{\partial}{\partial y_j} + A_j^{(1)} \right) \hat{g}^{jk} \left(-i \frac{\partial}{\partial y_k} + A_k^{(1)} \right) \hat{u}_1 \overline{\hat{v}_{1t}} \right. \\ \left. + \hat{u}_{1t} \sum_{j,k=1}^{n-1} \left(i \frac{\partial}{\partial y_j} + A_j^{(1)} \right) \hat{g}^{jk} \left(i \frac{\partial}{\partial y_k} + A_k^{(1)} \right) \overline{\hat{v}_1} \right] dy dt + \int_{X_{20}} V_1 \hat{u}_{1t} \overline{\hat{v}_1} dy dt + \int_{X_{20}} V_1 \hat{u}_1 \overline{\hat{v}_{1t}} dy dt \\ = \int_{X_{20}} \frac{\partial}{\partial t} \sum_{j,k=1}^{n-1} \hat{g}^{jk}(y) \left(-i \frac{\partial}{\partial y_k} + A_k^{(1)} \right) \overline{\left(-i \frac{\partial}{\partial y_j} + A_j^{(1)} \right)} \hat{u}_1 \overline{\hat{v}_1} dy dt + \int_{X_{20}} V_1 \frac{\partial}{\partial t} (\hat{u}_1 \overline{\hat{v}_1}) dy dt.$$

Let

$$(2.23) \quad t - y_n = s, \quad T - t - y_n = \tau,$$

i.e. $y_n = \frac{T-\tau-s}{2}$, $t = \frac{T-\tau+s}{2}$. Note that

$$u_s = \frac{1}{2}(u_t - u_{y_n}), \quad u_\tau = -\frac{1}{2}(u_t + u_{y_n})$$

and that $\tau = 0$ on Y_{20} . Combining (2.20), (2.21), (2.22) we get

$$(2.24) \quad 0 = (L_1 \hat{u}_1, \hat{v}_{1t}) + (\hat{u}_{1t}, L_1^* \hat{v}_1) = Q(\hat{u}_1, \hat{v}_1) - \Lambda_0(f, g),$$

where

$$(2.25) \quad Q(\hat{u}_1, \hat{v}_1) = \int_{Y_{20}} \frac{1}{2} \left[4 \hat{u}_{1s} \overline{\hat{v}_{1s}} + \sum_{j,k=1}^{n-1} \hat{g}^{jk} \left(-i \frac{\partial}{\partial y_j} + A_j^{(1)} \right) \hat{u}_1 \overline{\left(-i \frac{\partial}{\partial y_k} + A_k^{(1)} \right)} \hat{v}_1 + V_1(y) \hat{u}_1 \overline{\hat{v}_1} \right] dy' ds,$$

$$(2.26) \quad \Lambda_0(f, g) = - \int_{\Delta_{20}} \hat{u}_{1y_n} \overline{\hat{v}_{1t}} + \hat{u}_{1t} \overline{\hat{v}_{1y_n}} dy' dt.$$

Note that $u_{1t}|_{\Delta_{20}} = f_t$, $\hat{v}_{1t}|_{\Delta_{20}} = g_{1t}$, $\hat{u}_{1y_n}|_{\Delta_{20}} = \Lambda^{(1)} f$, $v_{1y_n}|_{\Delta_{20}} = \Lambda_*^{(1)} g$, where $\Lambda^{(1)}$ is the D-to-N operator on Δ_{20} .

Therefore $\Lambda_0(f, g)$ is known for all f, g in $H_0^1(\Delta_{20})$ if we know $\Lambda^{(1)}$. Here $H_0^1(\Delta_{js_0})$ is the closure in $H^1(\Delta_{js_0})$ of smooth functions equal to zero on $\partial\Delta_{js_0} \setminus \{t = T\}$, $0 \leq s_0 < T$, $j = 1, 2$.

For the convenience we shall often use the notation u^f, v^g where $L_1 u^f = 0$ and $L_1^* v^g = 0$ in X_{20} to emphasize the dependence of u^f and v^g on f and g respectively. Note that $Q(\hat{u}_1, \hat{v}_1)$ is a bounded nonsymmetric bilinear form on $H_0^1(Y_{20})$ where $H_0^1(Y_{js_0})$ consists of functions in $H^1(Y_{js_0})$ equal to zero on $\partial Y_{js_0} \setminus \{t = T\}$, $0 \leq s_0 < T$, $1 \leq j \leq 2$.

Note that for T small we have for any $\hat{u}_1 \in H_0^1(Y_{20})$

$$(2.27) \quad \Re Q(\hat{u}_1, \hat{u}_1) \geq C \|\hat{u}_1\|_1^2,$$

where $\|\cdot\|_1$ is the norm in $H^1(Y_{20})$. Also we have

$$(2.28) \quad |Q(\hat{u}_1, \hat{v}_1)| = |\Lambda_0(f, g)| \leq C \|f\|_1 \|g\|_1,$$

where $\|\cdot\|_1$ is the norm in $H^1(\Delta_{20})$.

Note that on the plane $\tau = 0$ we have $\Gamma \times [s_0, T] \subset \overline{Y_{1s_0}} \subset R_{s_0} \subset \overline{Y_{2s_0}}$ where $R_{s_0} = \Gamma^{(1)} \times [s_0, T]$.

Lemma 2.2. *For any $w \in H_0^1(R_{s_0}) \subset H_0^1(Y_{2s_0})$ there exists a sequence $\{u^{f_n}\}$, $f_n \in H_0^1(\Delta_{2s_0})$ that converges to w in $H_0^1(Y_{2s_0})$.*

We shall postpone the proof of Lemma 2.2 until the section 3.

Let the span of u_j , $j \geq 1$, be dense in $H_0^1(Y_{2s_0})$.

Lemma 2.3. *(c.f. [GLT]) For any $u \in H_0^1(Y_{20})$ there exists $u_0 \in H_0^1(Y_{2s_0})$ such that $Q(u_0, v') = Q(u, v')$, $\forall v' \in H_0^1(Y_{2s_0})$.*

Proof: Take $u^{(N)} = \sum_{j=1}^N c_{jN} u_j$ and determine c_{jN} from the linear system

$$(2.29) \quad Q(u^{(N)}, u_k) = Q(u, u_k), \quad 1 \leq k < N,$$

i.e.

$$(2.30) \quad \sum_{j=1}^N c_{jN} Q(u_j, u_k) = Q(u, u_k), \quad 1 \leq k \leq N.$$

Multiplying (2.29) by $\overline{c_{kN}}$ and adding, we get that

$$(2.31) \quad Q(u^{(N)}, u^{(N)}) = Q(u, u^{(N)}).$$

Therefore

$$C_1 \|u^{(N)}\|_1^2 \leq \Re Q(u^{(N)}, u^{(N)}) \leq C \|u\|_1 \|u^{(N)}\|_1.$$

Therefore system (2.29) has a unique solution and $\|u^{(N)}\|_1 \leq C_2 \|u\|_1$ for all N . Since the sequence $\|u^{(N)}\|_1$ is bounded it is weakly compact in $H_0^1(Y_{2s_0})$. Therefore there exists a subsequence $u^{(N_k)}$ such that $u^{(N_k)} \rightharpoonup u_0 \in H_0^1(Y_{2t_0})$ weakly when $N_k \rightarrow \infty$. Passing to the limit in (2.29) we get

$$(2.32) \quad Q(u_0, u_k) = Q(u, u_k), \quad \forall k.$$

We used that Q is continuous bilinear form in $H_0^1(Y_{2s_0})$. Since the span of $\{u_k\}$ is dense we get $Q(u_0, v') = Q(u, v')$, $\forall v' \in H_0^1(Y_{2s_0})$. Note that any subsequence of $\{u^{(N)}\}$ has a subsequence that converges to u_0 since (2.27) implies that there is a unique u_0 satisfying (2.32). Therefore the whole sequence $u^{(N)}$ converges weakly to u_0 . Note that $u^{(N)}$ converges also strongly in $H_0^1(Y_{2s_0})$ to u_0 since

$$Q(u_0 - u^{(N)}, u_0 - u^{(N)}) = Q(u_0, u_0) - Q(u_0, u^{(N)}) - Q(u^{(N)}, u_0) + Q(u^{(N)}, u^{(N)}).$$

Since $Q(u^{(N)}, u^{(N)}) = Q(u, u^{(N)})$ and $Q(u_0, u_0) = Q(u, u_0)$ we get that $Q(u_0 - u^{(N)}, u_0 - u^{(N)}) \rightarrow 0$ when $N \rightarrow \infty$. $\square$

Note that Lemma 2.3 remains true when $Q(u, v')$ is replaced by any linear continuous functional $\Phi(v')$ on $H_0^1(Y_{2s_0})$.

Applying the Green's formula in X_{20} we get, similarly to (2.24):

$$(2.33) \quad \begin{aligned} 0 &= \int_{X_{20}} (L_1 u \bar{v} - u \bar{L}_1^* v) dy dt \\ &= \int_{Y_{20}} (u_s \bar{v} - u \bar{v}_s) ds dy' - \int_{\Delta_{20}} (u_{y_n} \bar{v} - u \bar{v}_{y_n}) dy' dt. \end{aligned}$$

We used in (2.33) that $u = v = 0$ on Z_{20} . Integrating by parts we get

$$(2.34) \quad - \int_{Y_{20}} u \bar{v}_s ds dy' = - \int_{\gamma_{20}} u(y', 0, T) \overline{v(y', 0, T)} dy' + \int_{Y_{20}} u_s \bar{v} ds dy'.$$

Let

$$(2.35) \quad A(u, v) = 2 \int_{Y_{20}} u_s \bar{v} ds dy'.$$

Since $u(y', 0, T) = f(y', T)$ and $v(y', 0, T) = g(y', T)$ we get from (2.33) and (2.34) that $A(u, v)$ is determined by the boundary data.

Fix f, g in $H_0^1(\Delta_{10})$ and smooth. Let $s_0 \in [0, T)$ be arbitrary. We shall show that there exists a unique $u_0 \in H_0^1(R_{s_0})$ such that

$$(2.36) \quad A(u_0, v') = A(u^f, v'), \quad \forall v' \in H_0^1(Y_{2s_0}).$$

Note that $\overline{Y_{10}} \cap \{s \geq s_0\} \subset R_{s_0}$. Denote by $w(s, y')$ a function equal to $u^f|_{s=s_0}$ in R_{s_0} and equal to zero in $Y_{2s_0} \setminus R_{s_0}$. Then $\frac{\partial w}{\partial s} = 0$ in Y_{2s_0} , $u_0 = u^f - w \in H_0^1(R_{s_0})$ and $\frac{\partial u^f}{\partial s} = \frac{\partial u_0}{\partial s}$ in Y_{2s_0} . Therefore (2.36) holds.

We shall show that $A(u_0, v^g)$ is uniquely determined by the D-to-N operator.

Lemma 2.4. *Let $L_1^{(i)}$, $i = 1, 2$, be two formally self-adjoint operators in $X_{20}^{(i)}$, $i = 1, 2$, such that the corresponding D-to-N operators $\Lambda_1^{(i)}$, $i = 1, 2$, are equal on Δ_{2s_0} . Then*

$$C_1 \|u_1^f\|_{1, Y_{2s_0}^{(1)}} \leq \|u_2^f\|_{1, Y_{2s_0}^{(2)}} \leq C_2 \|u_1^f\|_{1, Y_{2s_0}^{(1)}}$$

for all $f \in H_0^1(\Delta_{2s_0})$.

Here $L_1^{(i)} u_i^f = 0$ in $X_{20}^{(i)}$.

We shall show first that $\Delta_{2s_0}^{(1)} = \Delta_{2s_0}^{(2)}$. Denote by $\Delta^{(i)}(0, T)$ the intersection of the domain of influence $D(\Delta_{10})$ of $L_1^{(i)}$ with $y_n = 0, i = 1, 2$. Note that $\Delta^{(i)}(0, T) = \overline{(\cup_f \text{supp } u_i^f)} \cap \{y_n = 0\}$ where the union is taken over all $f \in H_0^1(\Delta_{10})$. Denote $\tilde{\Delta}^{(i)} = (\Gamma \times [0, T]) \cup G_i$ where G_i is the closure of the union over all $f \in H_0^1(\Delta_{10})$ of $\text{supp } \Lambda^{(i)} f, i = 1, 2$. It is clear that $\tilde{\Delta}^{(i)} \subset \Delta^{(i)}(0, T), i = 1, 2$. The inclusion $\Delta^{(i)}(0, T) \subset \tilde{\Delta}^{(i)}$ follows from the local uniqueness of the Cauchy problem (see [T]). Since $\Lambda^{(1)} = \Lambda^{(2)}$ we have $\tilde{\Delta}^{(1)} = \tilde{\Delta}^{(2)}$. Therefore $\Delta^{(1)}(0, T) = \Delta^{(2)}(0, T)$. In particular, $\Gamma_1^{(1)} = \Gamma_2^{(1)}$ and therefore $\Delta_{2s_0}^{(1)} = \Delta_{2s_0}^{(2)}$. Now we shall proceed with the proof of Lemma 2.4.

Consider the identity (2.24) for $L_1^{(1)}$ and $L_1^{(2)}$ respectively assuming that $(L_1^{(i)})^* = L_1^{(i)}$ and $u_i^f = v_i^f, i = 1, 2$. Since $\Lambda_0^{(1)}(f, f) = \Lambda_0^{(2)}(f, f)$ we have that $Q^{(1)}(u_1^f, u_1^f) = Q^{(2)}(u_2^f, u_2^f)$ where the quadratic form $Q^{(i)}$ corresponds to $L_1^{(i)}$. Since T is small $Q^{(i)}(u_i^f, u_i^f)$ is equivalent to $\|u_i^f\|_{1, Y_{2s_0}^{(i)}}^2$. $\square$

We shall show that Lemma 2.4 implies that

$$(2.37) \quad A^{(1)}(u_0^{(1)}, v_1^g) = A^{(2)}(u_0^{(2)}, v_2^g)$$

where $A^{(i)}$ and $u_0^{(i)}$ correspond to $L_1^{(i)}$, $i = 1, 2$. It follows from (2.33), (2.34) that $A^{(1)}(u_1^{f'}, v_1^{g'}) = A^{(2)}(u_2^{f'}, v_2^{g'})$ for any $f' \in H_0^1(\Delta_{2s_0}), g' \in H_0^1(\Delta_{20})$.

By Lemma 2.2 there exists a sequence $f_n \in H_0^1(\Delta_{2s_0})$ such that $\|u_0^{(1)} - u_1^{f_n}\|_{1,Y_{2s_0}^{(1)}} \rightarrow 0$ when $n \rightarrow \infty$. By Lemma 2.4 the sequence $\{u_2^{f_n}\}$ is also convergent in $H_0^1(Y_{2s_0}^{(2)})$ to some $w^{(2)} \in H_0^1(Y_{2s_0}^{(2)})$. Since $A^{(i)}(u_i^{f'}, v_i^{g'})$ are continuous functionals on $H_0^1(Y_{2s_0}^{(i)})$, $i = 1, 2$, we get passing to the limit that

$$(2.38) \quad A^{(1)}(u_0^{(1)}, v_1^{g'}) = A^{(2)}(w^{(2)}, v_2^{g'}).$$

In (2.38) g' is arbitrary in $H_0^1(\Delta_{20})$. Take $g' \in H_0^1(\Delta_{2s_0})$. Then (2.36) implies that

$$(2.39) \quad A^{(1)}(u_0^{(1)}, v_1^{g'}) = A^{(2)}(u_0^{(2)}, v_2^{g'}).$$

Comparing (2.38) and (2.39) we get $A^{(2)}(u_0^{(2)}, v_2^{g'}) = A^{(2)}(w^{(2)}, v_2^{g'})$. By Lemma 2.2 $\{v_2^{g'}\}, g' \in H_0^1(\Delta_{2s_0})$ are dense in $H_0^1(R_{s_0})$. Therefore $u_0^{(2)} = w^{(2)}$ in R_{s_0} . We assume that $v^g \in H_0^1(Y_{10})$. Since $Y_{10} \cap \{s \geq s_0\} \subset R_{s_0}$ we get that $A^{(2)}(w^{(2)}, v^g) = A^{(2)}(u_0^{(2)}, v^g)$. Therefore (2.37) holds, i.e. $A(u_0, v^g)$ is uniquely determined by the D-to-N operator.

Remark 2.1 In this remark we shall show that in the self-adjoint case $A(u_0, v^g)$ can be recovered constructively from the boundary data. For any $\varepsilon > 0$ $Q_\varepsilon(u^f, v^g) = \varepsilon Q(u^f, v^g) + A(u^f, v^g)$ is determined by the D-to-N operator. Let $g_j, j \geq 1$ be a dense set in $H_0^1(\Delta_{2s_0})$. Denote by $\overline{H}$ the closure of the span of $\{v^{g_j}\}$ in $H_0^1(Y_{2s_0})$. Since $\Re A(v', v') \geq 0$ (see (2.35)) we get that $\Re Q_\varepsilon(v', v') \geq C\varepsilon \|v'\|_1^2$ for any $v' \in H_0^1(Y_{2s_0})$. It follows from (2.30) with Q replaced by Q_ε , u_j replaced by v^{g_j} and c_{jN} replaced by $c_{jN\varepsilon}$ that $c_{jN\varepsilon}$ are determined by the D-to-N operator. Repeating the proof of Lemma 2.3 we get that $u_\varepsilon^{(N)} = \sum_{j=1}^N c_{jN\varepsilon} v^{g_j}$ converges in $H_0^1(Y_{2s_0})$ to $u_\varepsilon \in \overline{H}$ such that $Q_\varepsilon(u_\varepsilon, v') = Q_\varepsilon(u^f, v')$, $\forall v' \in \overline{H}$. Since $Q_\varepsilon(u_\varepsilon^{(N)}, v^g)$ is determined by the D-to-N operator the limit $Q_\varepsilon(u_\varepsilon, v^g) = \lim_{N \rightarrow \infty} Q_\varepsilon(u_\varepsilon^{(N)}, v^g)$ is also determined by the D-to-N operator. Here $v^g \in H_0^1(Y_{10})$. Denote $w_\varepsilon = u_\varepsilon - u_0$, where u_0 is the same as in (2.36), $u_0 \in H_0^1(R_{s_0})$. We have, using (2.36), that

$$\varepsilon Q(w_\varepsilon, v') + A(w_\varepsilon, v') = \varepsilon Q(u^f - u_0, v').$$

Take $v' = w_\varepsilon$. Then

$$C\varepsilon \|w_\varepsilon\|_1^2 \leq C\varepsilon \|u^f - u_0\|_1 \|w_\varepsilon\|_1.$$

Therefore $\|w_\varepsilon\|_1 \leq C\|u^f - u_0\|_1$ for all ε . As in the proof of Lemma 2.3 we get that a sequence w_{ε_k} converges weakly in $\overline{H}$ to some $w \in \overline{H}$ and $(w_s, v') = 0 \quad \forall v' \in \overline{H}$. Since $\overline{H} \supset H_0^1(R_{s_0})$ we get that $w = 0$ in R_{s_0} . Since $R_{s_0} \supset Y_{10} \cap \{s \geq s_0\}$ we have that $(w_s, v^g) = 0$. Therefore $Q_\varepsilon(u_\varepsilon, v^g) \rightarrow A(u_0, v^g)$. Since this is true for any sequence ε_k we have $A(u_0, v^g) = \lim_{\varepsilon \rightarrow 0} Q_\varepsilon(u_\varepsilon, v^g)$, i.e. $A(u_0, v^g)$ is determined by the D-to-N operator on Δ_{20} . $\square$

Denote

$$A_1(u^f, v^g) = A(u^f, v^g) - A(u_0, v^g).$$

Then $A_1(u^f, v^g)$ is also determined by the D-to-N operator and

$$(2.40) \quad A_1(u^f, v^g) = 2 \int_{Y_{10} \cap \{s \leq s_0\}} \frac{\partial u^f}{\partial s} \overline{v^g} ds dy',$$

since $u_s^f - u_{0s} = 0$ when $s \geq s_0$, $u_0 = 0$ when $s \leq s_0$.

Now we shall construct a geometric optics solution $\hat{u}(y)$ of $L_1 \hat{u} = 0$ such that $\hat{u}(y, 0) = \hat{u}_t(y, 0) = 0$ for $y_n > 0$ and substitute it in (2.40) to recover v^g . We are looking for $\hat{u}$ in the form:

$$(2.41) \quad \hat{u} = u_N + u^{(N+1)},$$

where

$$u_N = e^{ik(s-s_0)} \sum_{p=0}^N \frac{1}{(ik)^p} a_p(s, \tau, y').$$

Substituting $\hat{u}$ in $L_1 \hat{u} = (-4 \frac{\partial^2}{\partial s \partial \tau} + L'_1) \hat{u} = 0$ we get the transport equations for a_p :

$$\frac{\partial a_0}{\partial \tau} = 0, \quad 4 \frac{\partial a_p}{\partial \tau} = L_1 a_{p-1}, \quad p \geq 1.$$

Choose $a_0(s, y') = \chi_1(s) \chi_2(y')$, where $\chi_1(s) \in C_0^\infty(\mathbf{R}^1)$, $\chi_1(s) = 1$ for $|s - s_0| < \delta$, $\chi_1(s) = 0$ for $|s - s_0| > 2\delta$, $\chi_2(y') = \frac{1}{\varepsilon^{n-1}} \chi_0(\frac{y' - y'_0}{\varepsilon})$, $\chi_0(y') \in C_0^\infty(\mathbf{R}^{n-1})$, $\chi_0(y') = 0$ for $|y'| > \delta$, $\int_{\mathbf{R}^{n-1}} \chi_0(y') dy' = 1$, δ is small, $y'_0 \in \Gamma$.

We define $a_p = \frac{1}{4} \int_{T-s}^\tau (L_1 a_{p-1}) d\tau'$, $1 \leq p \leq N$, and $u^{(N+1)}$ as the solution of

$$\begin{aligned} L_1 u^{(N+1)} &= -\frac{1}{4^N (ik)^N} (L_1 a_N) e^{ik(s-s_0)}, \\ u^{(N+1)} &= u_t^{(N+1)} = 0 \quad \text{when } t = 0, \\ u^{(N+1)} &= 0 \quad \text{when } y_n = 0. \end{aligned}$$

Note that $\text{supp } u_N$ is contained in a small neighborhood of the line $\{s = s_0, y' = y'_0\}$. Therefore $\text{supp } (u_N + u^{(N+1)}) \cap \{y_n = 0\} \subset \Delta_{10}$.

Substitute (2.41) into (2.40). Note that the principal term in k has the form

$$ik \int_{s < s_0} e^{ik(s-s_0)} \chi_1(s) \chi_2(y') \overline{v^g(y', s)} dy' ds,$$

where $v^g(y', s)$ is $v^g(y', s, \tau)$ for $\tau = 0$. Integrating by parts in s and taking the limit when $k \rightarrow \infty$ we get that the boundary data determine

$$(2.42) \quad \int_{\mathbf{R}^{n-1}} \chi_2(y') \overline{v^g(y', s_0)} dy'.$$

Taking limit in (2.42) when $\varepsilon \rightarrow 0$ we can recover $v^g(y'_0, s_0)$, $y'_0 \in \Gamma$, $0 < s_0 < T$, $\tau = 0$. Changing T to $T - \tau'$, $0 < \tau' < T$, we can analogously recover $v^g(y', y_n, t)$ for $y' \in \Gamma$, $0 \leq y_n \leq \frac{T}{2}$, $y_n \leq t \leq T - y_n$. In particular we determine $\hat{v}^g(y, t)$ and its time derivatives for $t = \frac{T}{2}$ and $y \in \mathcal{D}_{\frac{T}{2}}$ where $\mathcal{D}_{\frac{T}{2}} = \Gamma \times [0, \frac{T}{2}]$.

It is known (see [B1]) that $\hat{v}^g(y, \frac{T}{2})$, $g \in C_0^\infty(\Delta_{10})$ are dense in $H^m(\mathcal{D}_{\frac{T}{2}})$ for any $m \geq 0$. Since $L_1 v^g = 0$ we have

$$(2.43) \quad \begin{aligned} \frac{\partial^2}{\partial t^2} \hat{v}^g(y, t) - \frac{\partial^2 \hat{v}^g}{\partial y_n^2} - \sum_{j,k=1}^{n-1} \hat{g}^{jk}(y) \frac{\partial^2 \hat{v}^g}{\partial y_j \partial y_k} \\ - \sum_{j=1}^{n-1} B_j \frac{\partial^2 \hat{v}^g}{\partial y_j} + C(y) \hat{v}^g = 0, \end{aligned}$$

where B_j , $1 \leq j \leq n-1$, C depend on $\hat{g}^{jk}, A_j^{(1)}, V_1$ (see (2.15)).

Fix any $y = y_0 \in \mathcal{D}_{\frac{T}{2}}$ and let $t = \frac{T}{2}$. We can consider (2.43) as a linear system with unknowns $\hat{g}^{jk}(y_0)$, $B_j(y_0)$, $C(y_0)$, $g \in C_0^\infty(\Delta_{10})$ is arbitrary. If the rank of (2.43) is not maximal then there exist constants $\alpha_{jk}, \alpha_j, \alpha_0, \alpha_{jk} = \alpha_{kj}$, not all equal to zero, such that

$$(2.44) \quad \sum_{j,k=1}^{n-1} \alpha_{jk} \frac{\partial^2 \hat{v}^g(y_0, \frac{T}{2})}{\partial y_j \partial y_k} + \sum_{j=1}^{n-1} \alpha_j \frac{\partial \hat{v}^g(y_0, \frac{T}{2})}{\partial y_j} + \alpha_0 \hat{v}^g(y_0, \frac{T}{2}) = 0$$

for all $\hat{v}^g(y, t)$, $g \in C_0^\infty(\Delta_{10})$. Then we have a contradiction since $\{\hat{v}^g\}$ are dense in $H_0^N(\mathcal{D}_{\frac{T}{2}})$, $N \geq \frac{n}{2} + 2$. Therefore the system (2.43) has the

maximal rank and $\hat{g}^{jk}(y_0), B_j(y_0), C(y_0)$ are uniquely determined by (2.43). In particular, we recover $\|\hat{g}^{jk}(y_0)\|$ in $\mathcal{D}_{\frac{T}{2}}$. Knowing the metric and $B_j(y_0)$ we recover $A_j^{(1)}(y_0)$, $1 \leq j \leq n-1$. Finally knowing $\hat{g}^{jk}, A_j^{(1)}$ and $C(y_0)$ in $\mathcal{D}_{\frac{T}{2}}$ we can recover $\hat{V}(y_0)$. Lemma 2.1 is proven. $\square$

Remark 2.2 We shall show that the D-to-N operator on Δ_{20} determines the metric tensor and (2.11) on Δ_{20} . Consider the cotangent space T_0^* of Δ_{20} . Let ξ_0 be the dual variable to the time variable t . Therefore the points of T_0^* have the form (y', t, ξ', ξ_0) . The region $T_E^* \subset T_0^*$ where $|\xi_0| < \varepsilon|\xi'|$, ε is small, is a part of the "elliptic" region of T_0^* (see, for example, [H] or [E2]) where microlocally L behaves as an elliptic operator. Therefore in the region T_E^* we can find the parametrix for the D-to-N operator Λ as in the elliptic case (see, for example, [LU] or [E5], pp 54-55). Therefore we can recover the full symbol of Λ in T_E^* , in particular, the principal symbol of Λ in semigeodesic coordinates:

$$i \left(\sum_{j,k=1}^{n-1} g^{jk}(y', 0) \xi_j \xi_k - \xi_0^2 \right)^{\frac{1}{2}}$$

It follows from [LU], formula (1.8), (see also [E5]), that one can recover $\frac{\hat{g}_{yn}(y', 0)}{\hat{g}(y', 0)}$. $\square$

3 The conclusion of the proof of Lemma 2.1.

We shall start with the proof of Lemma 2.2. We shall show first that it is enough to prove that the set $\{u^f\}$ where $f \in C_0^\infty(\Delta_{2s_0})$ is dense in $\overset{\circ}{H}^1(R_{s_0})$ where $\overset{\circ}{H}^1(Y_{2s_0})$ is the closure of $C_0^\infty(Y_{2s_0})$ in the $H^1(Y_{2s_0})$ norm and $\overset{\circ}{H}^1(R_{s_0})$ is defined analogously. Suppose there exists $v \in H_0^1(R_{s_0})$ that does not belong to the closure $\overline{H}$ of $\{v^f\}, g \in H_0^1(\Delta_{2s_0})$. Then there exists $w \in H_0^1(Y_{2s_0})$ such that $(w, u^f)_1 = 0, \forall f \in H_0^1(\Delta_{2s_0})$, and $(w, v)_1 = 1$, where $(v, w)_1$ is the inner product in $H^1(Y_{2s_0})$.

Take any $u_1 \in C_0^\infty(R_{s_0})$. Since by the assumption $\{u^f\}, f \in C_0^\infty(\Delta_{2s_0})$ are dense in $\overset{\circ}{H}^1(R_{s_0})$ we get that $(w, u_1) = 0$. Integrating by parts we get $\int_{R_{s_0}} (-\Delta w + w) \overline{u_1} dy ds = 0$. Since $u_1 \in C_0^\infty(R_{s_0})$ is arbitrary we get $-\Delta w + w = 0$ in R_{s_0} . Fix $y_1 \in \Gamma^{(1)}$ and consider $f \in H_0^1(\Delta_{2s_0})$ having support in $\{|y' - y_1| < \varepsilon\} \times (T - \varepsilon, T]$, where $\varepsilon > 0$ is small. Then $\text{supp } u^f \subset R_{s_0}$.

Again integrating by parts we get

$$\begin{aligned} 0 &= (w, u^f)_1 = \int_{R_{s_0}} (-\Delta w + w) \overline{u^f} dy' ds \\ &+ \int_{\Gamma^{(1)}} \frac{\partial w}{\partial s} \overline{u^f}(y', T) dy' = \int_{\Gamma^{(1)}} \frac{\partial w}{\partial s} \overline{u^f}(y', T) dy'. \end{aligned}$$

Since $u^f = f(y', T)$ when $t = T$ and $f(y', T)$ can be chosen arbitrary near y_1 we get that $\frac{\partial w}{\partial s} = 0$ for $t = T, |y' - y_1| < \varepsilon$. Analogously we can prove that $\frac{\partial w}{\partial s} = 0$ on $\Gamma^{(1)} \times \{t = T\}$. Then $(w, v)_1 = \int_{R_{s_0}} (-\Delta w + w) \overline{v} ds dy' + \int_{\Gamma^{(1)}} \frac{\partial w}{\partial s} \overline{v} dy' = 0$ and this contradicts the assumption that $(w, v)_1 = 1$. $\square$

To complete the proof of Lemma 2.2 we need two more lemmas.

Denote by Δ_1 a domain in $\mathbf{R}^{n+1}$ bounded by three planes: $\Gamma_2 = \{\tau = T - t - y_n = 0, 0 \leq y_n \leq \frac{T}{2}\}$, $\Gamma_3 = \{s = t - y_n = 0, \frac{T}{2} \leq y_n \leq T\}$, $\Gamma_4 = \{t = T, 0 \leq y_n \leq T\}$. Let $\mathcal{H} = H^1(\Gamma_4) \times L_2(\Gamma_4)$ and let $\mathcal{H}_1$ be the space of pairs $\{\varphi, \psi\}$, $\varphi \in H^1(\Gamma_2)$, $\psi \in H^1(\Gamma_3)$, $\varphi = 0$ when $t = T$, $\psi = 0$ when $y_n = T$, $\varphi = \psi$ when $t = \frac{T}{2}$, with the norm $(\|\varphi\|_{1,\Gamma_2}^2 + \|\psi\|_{1,\Gamma_3}^2)^{\frac{1}{2}}$. We shall consider functions with compact support in $y' \in \mathbf{R}^{n-1}$.

Lemma 3.1. *For any $\{v_0, v_1\} \in \mathcal{H}_1$ there exist $\{w_0, w_1\} \in \mathcal{H}$ and $u \in H^1(\Delta_1)$ such that $L_1 u = 0$ in Δ_1 , $u|_{\Gamma_2} = v_0$, $u|_{\Gamma_3} = v_1$, $u|_{\Gamma_4} = w_0$, $\frac{\partial u}{\partial t}|_{\Gamma_4} = w_1$.*

Proof: For any smooth u, v with compact supports in y' such that $L_1 u = 0$, $L_1^* v = 0$ in Δ_1 we have

$$(3.1) \quad 0 = (u_t, L_1^* v) + (L_1 u, v_t) = E(u, v) - Q(u, v) - Q_1(u, v),$$

where $Q(u, v)$ is similar to (2.25), $Q_1(u, v)$ is a bilinear form on Γ_3 of the form (2.25) with $u_s, \overline{v_s}$ replaced by $u_\tau, \overline{v_\tau}$, and

$$\begin{aligned} &E(u, v) \\ &= \int_{\Gamma_4} \left[u_t \overline{v_t} + u_{y_n} \overline{v_{y_n}} + \sum_{j,k=1}^{n-1} \hat{g}^{jk} \left(-i \frac{\partial}{\partial x_j} + A_j^{(1)} \right) u \overline{\left(-i \frac{\partial}{\partial x_k} + \overline{A_k^{(1)}} \right) v} + \hat{V}_1 u \overline{v} \right] dy. \end{aligned}$$

Consider $0 = (L_1 u, u_t) + (u_t, L_1 u)$ in Δ_1 . Integrate by parts as in (3.1). When L_1 is not self-adjoint we get, in addition to $Q(u, u)$, $Q_1(u, u)$ and

$E(u, u)$, an integral $E_1(u, u)$ over Δ_1 that satisfies the following estimate:

$$(3.2) \quad |E_1(u, u)| \leq C \int_{\Delta_1} (|u_t|^2 + \sum_{j=1}^n |u_{y_j}|^2 + |u|^2) dy dt.$$

Denote by $\Delta_{1,T'}$ the domain bounded by Γ_2 , Γ_3 and $\Gamma_{4,T'}$ where $\Gamma_{4,T'}$ is the plane $\{t = T'\}$, $T' \in [\frac{T}{2}, T]$. Let $(u, v)_{\Delta_{1,T'}}$ be the L_2 inner product over $\Delta_{1,T'}$. Then integrating by parts in $0 = (L_1 u, u_t)_{\Delta_{1,T'}} + (u_t, L_1 u)_{\Delta_{1,T'}}$ we get, analogously to (3.1), (3.2), with Δ_1 replaced by $\Delta_{1,T'}$:

$$(3.3) \quad \|u_t\|_{0,\Gamma_{4,T'}}^2 + \|u\|_{1,\Gamma_{4,T'}}^2 \leq C(\|u\|_{1,\Gamma_{2,T'}}^2 + \|u\|_{1,\Gamma_{3,T'}}^2 + \int_{\frac{T}{2}}^{T'} (\|u_t\|_{0,\Gamma_{4,t}}^2 + \|u\|_{1,\Gamma_{4,t}}^2) dt.$$

Here $\Gamma_{2,T'}$, $\Gamma_{3,T'}$ are parts of Γ_2 and Γ_3 where $t \leq T'$. Since T is small we get from (3.3) that

$$(3.4) \quad \max_{T' \in [\frac{T}{2}, T]} (\|u_t\|_{0,\Gamma_{4,T'}}^2 + \|u\|_{1,\Gamma_{4,T'}}^2) \leq C(\|u\|_{1,\Gamma_2}^2 + \|u\|_{1,\Gamma_3}^2).$$

For any $w_0, w_1 \in C_0^\infty(\Gamma_4)$ there exists a smooth solution of the Cauchy problem $L_1 u = 0$, $u(y, T) = w_0$, $\frac{\partial u(y, T)}{\partial t} = w_1$ in the domain $t < T$ (see, for example, [H]). In particular, $\{u|_{\Gamma_2}, u|_{\Gamma_3}\} \in \mathcal{H}_1$.

We shall show that the image of the map $\{w_0, w_1\} \rightarrow \{u|_{\Gamma_2}, u|_{\Gamma_3}\}$ where $\{w_0, w_1\} \in \mathcal{H}$ are smooth, $L_1 u = 0$, is dense in $\mathcal{H}_1$. Suppose there exists $\{\varphi, \psi\} \in \mathcal{H}_1$ such that $(v, \varphi)_1 + (v_1, \psi)_1 = 0$, $\forall \{v, v_1\} \in \mathcal{H}_1$, $v = u|_{\Gamma_2}$, $v_1 = u|_{\Gamma_3}$, $L_1 u = 0$, u is smooth.

As in the Lemma 2.3 (see the remark after the end of the proof of Lemma 2.3) one can find $\{\varphi_0, \psi_0\} \in \mathcal{H}_1$ such that $Q(\varphi_0, v) + Q_1(\psi_0, v_1) = (\varphi, v)_1 + (\psi, v_1)_1$ for any $\{v, v_1\} \in \mathcal{H}_1$. Therefore $Q(\varphi_0, v) + Q_1(\psi_0, v_1) = 0$, $\forall \{v, v_1\}$, $v = u|_{\Gamma_2}$, $v_1 = u|_{\Gamma_3}$, u is smooth, $L_1 u = 0$. Here $(\varphi, v)_1$ is the inner product in $H^1(\Gamma_j)$, $j = 2, 3$.

Extend $\varphi_0(s, y')$ by 0 for $s > T$ and extend $\psi_0(\tau, y')$ by 0 for $\tau < -T$. Let

$$b(y', y_n, t) = \varphi_0(s, y') + \psi_0(\tau, y') - \varphi_0(0, y').$$

Note that $\varphi_0(0, y') = \psi_0(0, y')$, $b|_{\Gamma_2} = \varphi_0(s, y')$, $b|_{\Gamma_3} = \psi_0(\tau, y')$. Let $\tilde{\Delta}_1$ be the region $\tau < 0, s > 0$. Note that $\tilde{\Delta}_1 \supset \Delta_1$. Consider $L_1^* b = (-4 \frac{\partial^2}{\partial s \partial \tau} + (L_1')^*) b$. We have $\frac{\partial^2 b}{\partial s \partial \tau} = 0$ in $\tilde{\Delta}_1$ and $(L_1')^* b \in H_{0,-1}$ in $\tilde{\Delta}_1$ where $H_{p,r'}$ is the Sobolev space of order p in all variables and of order r' in y' . Let $f = -L_1^* b$

in $\tilde{\Delta}_1$ and $f = 0$ otherwise. Then $f \in H_{0,-1}(\mathbf{R}^{n+1})$ and $f = 0$ for $t < \frac{T}{2}$. It follows from [E2], for example, that there exists $u_0 \in H_{1,-1}(\mathbf{R}^{n+1})$ with the weight $e^{-\sigma t}$ such that $L_1^* u_0 = f$ in $\mathbf{R}^{n+1}$, $u_0 = 0$ for $t < \frac{T}{2}$. Denote $u^{(0)} = b(y', y_n, t) + u_0$. Then $L_1^* u^{(0)} = 0$ in $\tilde{\Delta}_1$, $u^{(0)}|_{\Gamma_2} = \varphi_0$, $u^{(0)}|_{\Gamma_3} = \psi_0$. We used that $u_0 = 0$ on Γ_3 and Γ_2 by the finite domain of dependence property since $u_0 = 0$ for $t < \frac{T}{2}$.

Denote $u_{10} = u^{(0)}|_{t=T}$, $u_{20} = \frac{\partial u^{(0)}}{\partial t}|_{t=T}$. Applying Green formula (3.1) to $u^{(0)}$ and v , where v is smooth, $L_1 v = 0$, $v|_{\Gamma_4} \in C_0^\infty(\Gamma_4)$ we get

$$(3.5) \quad E(u^{(0)}, v) = Q(u^{(0)}, v) + Q_1(u^{(0)}, v) = 0.$$

Since $u^{(0)}|_{t=T}$ and $\frac{\partial u^{(0)}}{\partial t}|_{t=T}$ are distributions, the pairing in (3.5) is understood as an extension of the L_2 -inner product. Since $v|_{t=T}$ and $\frac{\partial v}{\partial t}|_{t=T}$ are arbitrary and smooth we get that $u^{(0)}|_{t=T} = \frac{\partial u^{(0)}}{\partial t}|_{t=T} = 0$. Then by the uniqueness of the Cauchy problem (see [H]) we get that $u^{(0)} = 0$ in Δ_1 . In particular, $u^{(0)}|_{\Gamma_2} = \varphi_0 = 0$, $u^{(0)}|_{\Gamma_3} = \psi_0 = 0$. It follows from (3.4) that the image of the map $\{w_0, w_1\} \in \mathcal{H} \rightarrow \{u|_{\Gamma_2}, u|_{\Gamma_3}\} \in \mathcal{H}_1$ is onto.

Therefore for any $\{v^{(0)}, w^{(0)}\} \in \mathcal{H}_1$ there exists $v^{(1)} \in \mathring{H}^1(\Gamma_4)$, $v^{(2)} \in L_2(\Gamma_4)$ and $v \in H_1(\Delta_1)$ such that

$$(3.6) \quad L_1 v = 0 \text{ in } \Delta_1, \quad v|_{\Gamma_2} = v^{(0)}, \quad v|_{\Gamma_3} = w^{(0)}, \quad v|_{t=T} = v^{(1)}, \quad \frac{\partial v}{\partial t}|_{t=T} = v^{(2)}.$$

Lemma 3.1 is proven. $\square$

Denote by Δ_2 the domain bounded by $t = T$, $y_n = 0$ and $t - y_n = 0$. Note that $\Delta_1 \subset \Delta_2$. There exists (see, for example, [H]) a unique solution $u \in H^1(\Delta_2)$ of $L_1 u = 0$ in Δ_2 with the initial conditions $u|_{t=T} = v^{(1)}$, $\frac{\partial u}{\partial t}|_{t=T} = v^{(2)}$ and the boundary condition $u|_{y_n=0} = 0$. Here $v^{(1)}$ and $v^{(2)}$ are the same as in (3.6). By the uniqueness of the Cauchy problem $u = v$ in $\Delta_1 \subset \Delta_2$, where v is the solution obtained in (3.6).

Therefore we proved the following lemma:

Lemma 3.2. *For any $v_0 \in \mathring{H}^1(Y_{20}) \subset \mathring{H}^1(\Gamma_2)$ there exists $u \in H^1(X_{20})$ such that $L_1 u = 0$ in X_{20} , $u|_{Y_{20}} = v_0$, $u|_{y_n=0} = 0$.*

Analogous result holds when X_{20} is replaced by X_{2s_0} and L_1 is replaced by L_1^* .

Finally we can finish the proof of Lemma 2.2. Suppose that the set H of $\{u^g\}$, $g \in C_0^\infty(\Delta_{2s_0})$ is not dense in $\mathring{H}^1(R_{s_0})$. Then there exists $w_0 \in \mathring{H}^1(Y_{2s_0})$ such that $(u^g, w_0)_1 = 0$, $\forall u^g$, $g \in C_0^\infty(\Delta_{2s_0})$ and $(w_0, v) = 1$ for some $v \in \mathring{H}^1(R_{s_0})$. By Lemma 2.3 one can find $w \in \mathring{H}^1(Y_{2s_0})$ such that $Q(u^g, w) = (u^g, w_0)_1$ for all u^g . By Lemma 3.2 there exists $u \in H^1(X_{2s_0})$ such that $L_1^* u = 0$ in X_{2s_0} , $u|_{y_n=0} = 0$, $u|_{Y_{2s_0}} = w$.

Applying the Green's formula (2.24) we get

$$0 = Q(u^g, w) = - \int_{\Delta_{2s_0}} \frac{\partial u^g}{\partial y_n} \overline{u_t} dy' dt - \int_{\Delta_{2s_0}} \frac{\partial g}{\partial t} \frac{\partial \overline{u}}{\partial y_n} dy' dt$$

since $u^g|_{y_n=0} = g$. Since $u|_{y_n=0} = 0$ we get that

$$\int_{\Delta_{2s_0}} \frac{\partial g}{\partial t} \frac{\partial \overline{u}}{\partial y_n} dy' dt = 0, \quad \forall g \in C_0^\infty(\Delta_{2s_0}).$$

Since g is arbitrary we get that $\frac{\partial}{\partial t} \frac{\partial u}{\partial y_n} = 0$ in $\Delta_{2s_0} = \Gamma^{(1)} \times (s_0, T]$. Therefore $\frac{\partial u}{\partial t}$ satisfies $L_1^* \frac{\partial u}{\partial t} = 0$ in X_{2s_0} , $\frac{\partial u}{\partial t} = \frac{\partial}{\partial y_n} \frac{\partial u}{\partial t} = 0$ on Δ_{2s_0} . By the unique continuation theorem (see [T]) we get that $\frac{\partial u}{\partial t} = 0$ in the double cone of influence of $\Gamma^{(1)} \times [s_0, T]$. Note that the intersection of this double cone with the plane $\tau = 0$ is R_{s_0} . Therefore $w = u|_{Y_{2s_0}}$ satisfies the elliptic equation $(4 \frac{\partial^2}{\partial s^2} + (L_1^*)^*) w = 0$ on R_{s_0} since $\frac{\partial^2 u}{\partial t^2} = 0$, $\frac{\partial^2 u}{\partial s^2} = \frac{1}{4} (\frac{\partial}{\partial t} - \frac{\partial}{\partial y_n})^2 u = \frac{1}{4} \frac{\partial^2 u}{\partial y_n^2}$.

Consider $Q(w, v)$. Note that $\text{supp } v \in \mathring{H}^1(R_{s_0})$. Integrating by parts and using that w satisfying the equation $4w_{ss} + (L_1^*)^* w = 0$ in R_{s_0} we get that $Q(w, v) = 0$ and this contradicts the condition that $Q(w, v) = 1$. Therefore H is dense in $\mathring{H}^1(R_{s_0})$. $\square$

Let $L^{(p)} u^{(p)} = 0$, $p = 1, 2$, be two hyperbolic equations of the form (1.1) in domains $\Omega^{(p)}$, $p = 1, 2$, respectively, satisfying the initial-boundary conditions (1.2). We assume that $\Gamma_0 \subset \partial\Omega^{(1)} \cap \partial\Omega^{(2)}$, $\text{supp } f \subset \Gamma_0 \times (0, T_0]$ and $\Lambda^{(1)} = \Lambda^{(2)}$ on $\Gamma_0 \times (0, T_0)$ where $\Lambda^{(p)}$ is the D-to-N operator corresponding to $L^{(p)}$, $p = 1, 2$.

Let $B \subset \Omega^{(1)} \cap \Omega^{(2)}$ be homeomorphic to a ball and such that the domains $\Omega_i = \Omega^{(i)} \setminus \overline{B}$, $i = 1, 2$, are smooth. Assume that $\partial B \cap \partial\Omega^{(i)} \subset \Gamma_0$, $i = 1, 2$, and connected. Let $\gamma^{(i)} = \partial\Omega^{(i)} \setminus \overline{\Gamma_0}$, $i = 1, 2$. Note that $\partial\Omega_1 \setminus \gamma^{(1)} = \partial\Omega_2 \setminus \gamma^{(2)} \stackrel{\text{def}}{=} \Gamma_1$.

We conclude this section with the following lemma that will be important in the global step (c.f. [KKL1], Lemma 9):

Lemma 3.3. *Let $L^{(1)} = L^{(2)}$ in $\overline{B}$ and let $\delta = \max_{x \in \overline{B}} d(x, \Gamma_0)$, where $d(x, \Gamma_0)$ is the distance in $\overline{B}$ from $x \in \overline{B}$ to Γ_0 . Let Λ_i be the D-to-N operators corresponding to $L^{(i)}$ in smaller domains $\Omega_i, i = 1, 2$. If $\Lambda^{(1)} = \Lambda^{(2)}$ on $\Gamma_0 \times (0, T_0)$ then $\Lambda_1 = \Lambda_2$ on $\Gamma_1 \times (\delta, T_0 - \delta)$.*

Proof: Let Ω be either $\Omega^{(1)}$ or $\Omega^{(2)}$. Denote by $\Omega_T = \Omega \times (-\infty, T)$, $\partial\Omega_T = \partial\Omega \times (-\infty, T)$. Let $H_s^+(\Omega_T)$ be the Sobolev space with norm $\|u\|_s$ where $u = 0$ for $t < 0$. The proof of Lemma 3.3 will be based on a variant of the Runge theorem.

Lemma 3.4. *Denote by $\tilde{\Omega} = \Omega \setminus \overline{B}$, $\gamma = \partial\Omega \setminus \overline{\Gamma_0}$. Let $u \in H_{1+s}^+(\tilde{\Omega}_T)$ be the solution of*

$$(3.7) \quad \begin{aligned} Pu &= 0, \quad (x, t) \in \tilde{\Omega}_T, \\ u|_{\partial\tilde{\Omega}_T} &= f, \quad u = 0 \text{ for } t < \delta, \end{aligned}$$

where P has the form (1.1), $f \in H_{s+1}^+(\partial\tilde{\Omega}_T)$, $f = 0$ on $\gamma_T = \gamma \times (-\infty, T)$, $f = 0$ for $t < \delta$, $s \leq 0$. Then there exists a sequence of smooth functions u_n in Ω_T such that

$$(3.8) \quad Pu_n = 0, \quad (x, t) \in \Omega_T, \quad u_n|_{\partial\Omega_T} = f_n,$$

where $u_n = 0$ for $t < 0$, $f_n = 0$ on γ_T and $u_n \rightarrow u$ when $n \rightarrow \infty$ in $H_s^+(\tilde{\Omega}_T)$.

Proof: Let Ω_ε be a smooth domain in $\mathbf{R}^n$ such that $\Omega_\varepsilon \supset \Omega$, $\partial\Omega_\varepsilon = \gamma \cup \gamma_\varepsilon$, and Γ_0 is inside of Ω_ε .

The following existence and uniqueness lemma is well known (see, for example, [E2]).

Lemma 3.5. *For any $h \in H_s^+(\Omega_T)$ and $f \in H_{s+1}^+(\partial\Omega_T)$, $s \geq 0$, there exists a unique $u \in H_{1+s}^+(\Omega_T)$ such that*

$$Pu = h \text{ in } \Omega_T, \quad u|_{\partial\Omega_T} = f.$$

Moreover, $\frac{\partial u}{\partial \nu}|_{\partial\Omega_T} \in H_s^+(\partial\Omega_T)$ and

$$\|u\|_{1+s} + \left\| \frac{\partial u}{\partial \nu} \right\|_s \leq C\|f\|_{s+1} + C\|h\|_s.$$

When $h = 0$ the same result is true for any $s \in \mathbf{R}$.

The last statement follows from the fact that when $Pu = 0$ and $s < 0$ the norm $\|u\|_{1+s}$ is equivalent near $\partial\Omega_T$ to the norm $\|u\|_{1,s}$ (c.f. [E2]). Here $\|u\|_{p,s}$ is the Sobolev norm of order p in all variables and of order s in (y', t) .

Denote by $E(x, t, y, \tau)$ the forward Green function for the domain Ω_ε . More precisely $E(x, t, y, \tau)$ satisfies

$$\begin{aligned} PE &= \delta(x - y)\delta(t - \tau), \quad (x, t) \in \Omega_\varepsilon \times (-\infty, \infty), \quad (y, \tau) \in \Omega_\varepsilon \times (-\infty, +\infty), \\ E &= 0 \quad \text{for } t < \tau, \\ E &= 0 \quad \text{when } (x, t) \in \partial\Omega_\varepsilon \times (-\infty, +\infty), \quad (y, \tau) \in \Omega_\varepsilon \times (-\infty, +\infty). \end{aligned}$$

The existence of $E(x, t, y, \tau)$ follows from the Hadamard construction (see [H]) and the Lemma 3.5.

Denote by E the operator with the kernel $E(x, t, y, \tau)$. Let $D_\varepsilon = (\Omega_\varepsilon \setminus \bar{\Omega}) \times (0, T + \delta)$. Denote by K the closure of $E\varphi$, $\forall \varphi \in C_0^\infty(D_\varepsilon)$, in the norm of $H_s^+(\tilde{\Omega}_T)$, $s \leq 0$. Let $K^\perp \subset \overset{\circ}{H}_{-s}^-(\tilde{\Omega} \times (0, T))$ be the "orthogonal" complement to K , i.e. $(f, g) = 0$ for all $f \in K$ iff $g \in K^\perp$. Here (f, g) is the extension of the L_2 scalar product in $\tilde{\Omega} \times (0, T)$ and $\overset{\circ}{H}_{-s}^-(\tilde{\Omega} \times (0, T))$ consists of functions that belong to $H_{-s}(\mathbf{R}^n \times (0, +\infty))$ after being extended by zero for $t > T$ and for $x \notin \tilde{\Omega}$. Note that $(f, g) = (lf, g_+)$ where $lf \in H_s(\mathbf{R}^n \times (-\infty, +\infty))$ and $g_+ \in H_{-s}(\mathbf{R}^n \times (-\infty, +\infty))$ are arbitrary extensions of f and g such that $g_+ = 0$ for $x \notin \tilde{\Omega}$, $g_+ = 0$ for $t > T$, $lf = 0$ for $t < 0$. Let $g \in K^\perp$ be arbitrary and let $\overline{w(y, \tau)} = \overline{E^*g_+}$ or, formally:

$$\overline{w(y, \tau)} = \int_{-\infty}^{\infty} \int_{\mathbf{R}^n} E(x, t, y, \tau) \overline{g_+(x, t)} dx dt.$$

We have $(E\varphi, g_+) = (\varphi, E^*g_+) = 0$ for any $\varphi \in C_0^\infty(D_\varepsilon)$. Therefore $w(y, \tau) = 0$ for $(y, \tau) \in D_\varepsilon$, $w(y, \tau)$ is the solution of the initial-boundary value problem

$$(3.9) \quad \begin{aligned} P^*w &= g_+(y, \tau), \quad (y, \tau) \in (\Omega_\varepsilon \times (0, +\infty)), \\ w(y, \tau) &= 0 \quad \text{for } \tau > T, \quad y \in \Omega_\varepsilon, \quad w|_{\partial\Omega_\varepsilon \times (0, +\infty)} = 0, \end{aligned}$$

and $\overline{E(x, t, y, \tau)}$ is the Schwartz kernel of the solution operator to (3.9). Here P^* is the adjoint to P .

It follows from Lemma 3.5 that $w \in H_{1-s}(\Omega_\varepsilon \times (0, +\infty))$, $w = 0$ for $t > T$, $w = 0$ on $\partial\Omega_\varepsilon \times (0, +\infty)$ and $\frac{\partial w}{\partial \nu}|_{\partial\Omega_\varepsilon \times (0, +\infty)} \in H_{-s}(\partial\Omega_\varepsilon \times (0, +\infty))$. Since $w(y, \tau) = 0$ in D_ε and $g_+ = 0$ in $(\Omega_\varepsilon \setminus \tilde{\Omega}) \times (0, +\infty)$ we get by the

uniqueness of the Cauchy problem (see [T]) that $w = 0$ in $(\Omega_\varepsilon \setminus \bar{\Omega}) \times (\delta, +\infty)$. Therefore w and $\frac{\partial w}{\partial \nu}$ are zero on $(\partial\tilde{\Omega} \setminus \gamma) \times (\delta, +\infty)$. Take any $u \in H_{1+s}^+(\tilde{\Omega}_T)$ that satisfies (3.7). We have $(u, g_+) = (u, P^*w)$. Using the Green's formula in $\tilde{\Omega} \times (-\infty, \infty)$ we get $(u, g_+) = (Pu, w) = 0$, i.e. $u \in K$. $\square$

Now we shall prove Lemma 3.3.

Let v_i , $i = 1, 2$, be smooth solutions of $L^{(i)}v_i = 0$ in $\Omega_i \times (-\infty, T_0 - \delta)$, $v_i = 0$ for $t < \delta$ and such that $v_i = f$ on $\Gamma_1 \times (-\infty, T_0 - \delta)$, $\text{supp } f \subset \Gamma_1 \times (-\infty, T_0 - \delta)$. Extend f smoothly from $\partial\Omega_i \times (-\infty, T_0 - \delta)$ to $\partial\Omega_i \times (-\infty, T_0)$, $\text{supp } f \subset \Gamma_1 \times (-\infty, T_0)$. Then extend v_i in Ω_{i,T_0} as solutions of $L^{(i)}v_i = 0$, $v_i = f$ on $\partial\Omega_{i,T_0}$. We shall show that $\Lambda_1 f = \Lambda_2 f$ on $\Gamma_1 \times (\delta, T_0 - \delta)$. By Lemma 3.4 there exists a sequence of smooth functions w_{n1} in $\Omega_{T_0}^{(1)}$ satisfying $L^{(1)}w_{n1} = 0$ such that $\|v_1 - w_{n1}\|_s \rightarrow 0$ when $n \rightarrow \infty$. Here $\|\cdot\|_s$ is the norm in $H_s(\Omega_{1,T_0})$. Denote $f_n = w_{n1}|_{\partial\Omega_{T_0}^{(1)}} \in H_{s+\frac{1}{2}}^+(\partial\Omega_{T_0}^{(1)})$. Note that $\text{supp } f_n \subset \Gamma_{0,T_0}$. By Lemma 3.5 there exists $w_{n2} \in H_{s+\frac{1}{2}}^+(\Omega_{T_0}^{(2)})$ such that $L^{(2)}w_{n2} = 0$ in $\Omega_{T_0}^{(2)}$, $w_{n2} = f_n$ on $\partial\Omega_{T_0}^{(2)}$, $f_n = 0$ on $\gamma_{T_0}^{(2)}$. Since $\Lambda^{(1)} = \Lambda^{(2)}$ on Γ_{0,T_0} we have $\Lambda^{(1)}f_n = \Lambda^{(2)}f_n$ on Γ_{0,T_0} . Since $L^{(1)} = L^{(2)}$ in B the uniqueness of the Cauchy problem (see [T]) implies that $w_{n1} = w_{n2}$ and $\frac{\partial w_{n1}}{\partial \nu} = \frac{\partial w_{n2}}{\partial \nu}$ on $\Gamma_1 \times (-\infty, T_0 - \delta)$.

It follows from $L^{(1)}w_{n1} = 0$ and $w_{n1} \in H_s^+(\Omega_{1,T_0})$ that $w_{n1} \in H_{2,s-2}^+$ near $\partial\Omega_{1,T_0}$ (c.f. [E2]). Therefore $w_{n1}|_{\partial\Omega_{1,T_0}} = f_{n1} \in H_{s-\frac{1}{2}}(\partial\Omega_{1,T_0})$, $\frac{\partial w_{n1}}{\partial \nu}|_{\partial\Omega_{1,T_0}} = g_{n1} \in H_{s-\frac{3}{2}}^+(\partial\Omega_{1,T})$ and $\|f_{n1} - f\|_{s-\frac{1}{2}} + \|g_{n1} - \frac{\partial v_1}{\partial \nu}\|_{s-\frac{3}{2}} \leq C\|w_{n1} - v_1\|_s \rightarrow 0$ when $n \rightarrow \infty$. Denote

$$\frac{\partial v_i}{\partial \nu}|_{\partial\Omega_i \times (\delta, T-\delta)} = g_i, \quad i = 1, 2, \quad f_{n2} = w_{n2}|_{\partial\Omega_2 \times (0, T-\delta)}, \quad g_{n2} = \frac{\partial w_{n2}}{\partial \nu}|_{\partial\Omega_2 \times (0, T-\delta)}.$$

Since $f_{n1} = f_{n2}$ on $\Gamma_1 \times (0, T - \delta)$ we have that $\|f_{n2} - f\|'_{s-\frac{1}{2}} \rightarrow 0$ when $n \rightarrow \infty$ where $\|\cdot\|'_s$ means the norm in $H_s^+(\partial\Omega_{2,T-\delta})$. Therefore applying Lemma 3.5 to $v_2 - w_{n2}$ we get that $\|g_2 - g_{n2}\|'_{s-\frac{3}{2}} \leq C\|f_{n2} - f\|'_{s-\frac{1}{2}} \rightarrow 0$ when $n \rightarrow \infty$. Since $g_{n1} = g_{n2}$ on $\Gamma_1 \times (0, T - \delta)$ we get that $g_1 = g_2$ on $\Gamma_1 \times (\delta, T - \delta)$. $\square$

Acknowledgements. Author expresses deep gratitude to Jim Ralston for his generous help. My special thanks to Slava Kurylev who saved me from embarrassing mistakes and gave many useful suggestions that greatly improved this paper.

References

- [B1] Belishev, M., 1997, Boundary control in reconstruction of manifolds and metrics (the BC method), *Inverse Problems* 13, R1-R45
- [B2] Belishev, M., 2002, How to see waves under the Earth's surface (the BC-method for geophysicists), *Ill -Posed and Inverse Problems*, 55-72 (S.Kabanikhin and V.Romanov (Eds), VSP)
- [E1] Eskin, G., 2004, Inverse boundary value problems in domains with several obstacles, *Inverse Problems* 20, 1497-1516
- [E2] Eskin, G., 1987, Mixed initial-boundary value problems for second order hyperbolic equations, *Comm. in PDE*, 12, 503-87
- [E3] Eskin, G., 1998, Inverse scattering problem in anisotropic media, *Commun. Math. Phys.* 199, 471-91
- [E4] Eskin, G., 2005, Inverse problems for Schrödinger equations with Yang-Mills potentials in domains with obstacles and the Aharonov-Bohm effect, *Journal of Physics Conference Series*, 12, 23-32
- [E5] Eskin, G., 2003, Inverse boundary value problems and the Aharonov-Bohm effect, *Inverse problems*, 19, 49-62
- [H] Hormander, L., 1985, *The Analysis of Linear Partial Differential Operators III* (Berlin: Springer)
- [GLT] Glowinski, R., Lions, J.L., Tremolieres, R., 1971, *Analyse numerique de inequations variationnelles* (Paris, Dunod)
- [I] Isakov, V., 1998, *Inverse problems for partial differential equations*, *Appl. Math. Studies*, vol. 127, Springer, 284 pp.
- [K] Kurylev, Y., 1993, Multi-dimensional inverse boundary problems by BC-mathod : groups of transformations and uniqueness results, *Math. Comput. Modelling* 18, 33-45
- [KK] Katchalov, A., Kurylev, Y., 1998, Multidimensional inverse problems with incomplete boundary spectral data, *Comm. Part. Diff. Eq.* 23, 55-95

- [KKL] Katchalov, A., Kurylev, Y., Lassas, M., 2001, Inverse boundary spectral problems (Boca Baton : Chapman&Hall)
- [KKL1] Katchalov, A., Kurylev, Y., Lassas, M., 2004, Energy measurements and equivalence of boundary data for inverse problems on noncompact manifolds, IMA Volumes, v.137, 183-214
- [KL1] Kurylev, Y. and Lassas, M., 2000, Hyperbolic inverse problems with data on a part of the boundary AMS/1P Stud. Adv. Math, 16, 259-272
- [LU] Lee, J. and Uhlmann, G., 1989, Determining anisotropic real-analytic conductivity by boundary measurements, Comm. Pure Appl. Math. 42, 1097-1112
- [T] Tataru, D., 1995, Unique continuation for solutions to PDE, Comm. in PDE 20, 855-84